

**DOI: 10.5152/etd.2018.0103**

**Manuscript Type:** Case Report

**Title:** De Novo Mutation in ATP7A Gene with Severe Menkes Disease

**Running Head:** Menkes Disease

**Authors:** Pembe Soylu Üstkoyuncu<sup>1</sup>, Ahmet Sami Güven<sup>2</sup>, Aslıhan Kiraz<sup>3</sup>, Ayşegül Yılmaz<sup>4</sup>, Şefika Elmas Bozdemir<sup>5</sup>, Songül Gökay<sup>1</sup>

**Institutions:**

<sup>1</sup>Department of Pediatric Nutrition and Metabolism, Health Sciences University, Kayseri Training and Research Hospital, Kayseri, Turkey

<sup>2</sup>Department of Pediatric Neurology, Health Sciences University, Kayseri Training and Research Hospital, Kayseri, Turkey

<sup>3</sup>Department of Genetic Clinic, Health Sciences University, Kayseri Training and Research Hospital, Kayseri, Turkey

<sup>4</sup>Department of Pediatric Genetic Clinic, Health Sciences University, Kayseri Training and Research Hospital, Kayseri, Turkey

<sup>5</sup>Department of Pediatric Infectious Disease, Health Sciences University, Kayseri Training and Research Hospital, Kayseri, Turkey

**Address for Correspondence:** Pembe Soylu Üstkoyuncu

Department of Pediatric Nutrition and Metabolism, Health Sciences University, Kayseri Training and Research Hospital, Kayseri, Turkey

e-mail: drpembesoylu@erciyes.edu.tr

Phone: 0352 336 88 88

**Submitted:** 08.9.2017

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as: Soylu Üstkoyuncu P, Güven AS, Kiraz A, Yılmaz A, Bozdemir ŞE, Gökay S. De Novo Mutation in ATP7A Gene with Severe Menkes Disease. Erciyes Med J DOI: 10.5152/etd.2018.0103

©Copyright 2018 by Erciyes University Faculty of Medicine - Available online at [www.erciyesmedj.com](http://www.erciyesmedj.com)

## **ABSTRACT**

Menkes disease (MD) is an X-linked neurodegenerative disorder, which occurs in early infancy, and is caused by the impairment of P-type ATPase.

An 8-month-old boy presented with seizure and difficulty of feeding. His hair was blond, thin, and weak. He had poor head control and could not sit. The microscopic appearance of the patient's hair was pili torti. Brain magnetic resonance imaging revealed diffuse cerebral and cerebellar atrophy, and vascular tortuosity was observed in both middle cerebral and vertebral basilar arteries in magnetic resonance angiography. Molecular genetic analysis was performed for suspected MD and a hemizygous mutation (p. G1118S [c.3352G>A]) was detected in ATP7A gene.

Although it is not specific for the disorder, microscopy of the hair allows early diagnosis when the differential diagnosis is broad or other tests are not conclusive. Since it is a fatal neurodegenerative disorder, genetic counseling must be provided to the family.

**Keywords:** ATP7A, De Novo Mutation, Menkes Disease

## **INTRODUCTION**

Menkes disease (MD) is an X-linked neurodegenerative disorder, which occurs in early infancy, and is caused by the impairment of P-type ATPase (ATP7A), which is essential for copper homeostasis. The responsible gene was mapped on Xq21.1. The disorder is characterized by hypotonia, feeding

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as: Soylu Üstkoyuncu P, Güven AS, Kiraz A, Yılmaz A, Bozdemir ŞE, Gökay S. De Novo Mutation in ATP7A Gene with Severe Menkes Disease. *Erciyes Med J* DOI: 10.5152/etd.2018.0103

difficulties, seizures, dysmorphic facial features, psychomotor retardation, and connective tissue anomalies (1).

Gastrostomy tube replacement for the management of feeding, antiepileptic drugs for seizures, and surgery for bladder diverticula and/or subdural hematomas are some treatment options for supportive care. It has been reported that copper histidine reduces seizure frequency when the treatment is started in the early period (2).

## **CASE REPORT**

An 8-month-old male infant presented with seizure and difficulty of feeding. His body weight was 8.2 kg (25%–50% percentile), height was 73 cm (50%–75% percentile), and the head circumference was 45 cm (25%–50% percentile). His hair was blond, thin, and weak (Figure 1a). He had poor head control and he could not sit.

The patient was the second child of a nonconsanguineous Turkish couple. Prenatal, obstetric, and family history were unremarkable. The parents also had a 2-year-old healthy daughter.

In laboratory investigations, complete blood count, blood glucose, liver enzymes, and creatine kinase levels were within normal limits.

He fixed his vision on one point; his activities reduced for a few minutes in his first days of hospitalization. Electroencephalography (EEG) revealed a sharp wave activity in the temporo-occipital and parieto-occipital regions of the left hemisphere also slowing of the ground rhythm in the posterior half of both hemispheres and temporal region of the left hemisphere. Due to the presence of hypomotor seizures, phenobarbital was started and seizures were controlled.

Brain magnetic resonance imaging (MRI) revealed diffuse cerebral and cerebellar atrophy; magnetic resonance angiography (MRA) showed vascular tortuosity in both the middle cerebral and vertebrobasilar arteries (Figure 2).

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as: Soylu Üstkoyuncu P, Güven AS, Kiraz A, Yılmaz A, Bozdemir ŞE, Gökay S. De Novo Mutation in ATP7A Gene with Severe Menkes Disease. *Erciyes Med J* DOI: 10.5152/etd.2018.0103

Metabolic studies were performed due to psychomotor retardation and seizures. Blood lactate, pyruvate, biotinidase activity, plasma and urine amino acid analysis, tandem mass spectroscopy, and urine organic acid analysis were normal.

The microscopic appearance of the patient's hair was pili torti (Figure 1b). Therefore, the patient was evaluated for MD. Serum ceruloplasmin level was 0.08 g/L (normal range [NR]: 0.31–0.91), and serum copper level was 22 g/dL (NR: 50–155).

Molecular genetic analysis was performed for suspected MD and a hemizygous mutation (p. G1118S [c.3352G>A]) was detected in the ATP7A gene (Figure 3a). The information about the risk of transmitting the ATP7A mutation was provided to the family. Mutation-targeted molecular testing was performed to proband and the females at risk by the Sanger sequencing method (ABI 3500). They have no mutation in the suspected gene region (Figure 3b). These results suggested that the mutation of the proband is a de novo mutation. Pedigree of the family is shown in Figure 3c.

The patient is 21 months old now. He is bedridden and has a significant feeding difficulty. He was hospitalized due to pulmonary infection several times.

## DISCUSSION

Although hypotonia, hypothermia, feeding difficulties, seizures, dysmorphic facial features, and cognitive and motor retardation are major clinical findings of MD, they are not specific. It can be misdiagnosed as many chronic neurological diseases.

Seizures are common in MD. Early onset seizures typically start with focal seizures and show a progression to epileptic spasms. The late forms of chronic seizures can appear as tonic seizures, myoclonic jerks, infantile spasms, multifocal epileptiform activity, or hypersarrhythmia on EEG. Rizk et al. (3) reported a patient with epilepsy partialis continua, and MRA of this patient showed relatively tortuous but patent intracranial vessels, with an appearance of a "hair pin" sign. Middle cerebral,

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as: Soylu Üstkoyuncu P, Güven AS, Kiraz A, Yılmaz A, Bozdemir ŞE, Gökay S. De Novo Mutation in ATP7A Gene with Severe Menkes Disease. *Erciyes Med J* DOI: 10.5152/etd.2018.0103

vertebral, and basilar arteries are elongated and tortuous in MRA, as in the present case. Our patient presented with hypomotor seizures and, which were controlled with phenobarbital.

Light-colored hair can be seen in phenylketonuria and trichorhexis nodosa; pili torti can be seen in ectodermal dysplasia, argininosuccinic aciduria, or biotinidase deficiency. Metabolic tests of our case were normal. Therefore, these inborn errors of metabolic diseases have been excluded.

Although microscopy of the hair allows early diagnosis when the differential diagnosis is broad and/or other tests are not conclusive, it is not specific for MD (4). Serum copper and ceruloplasmin levels are supportive diagnostic tests, but they are also not specific for MD.

To date, 274 different disease-causing variations of ATP7A have been reported, but some of them are not disease related. Splice site nonsense mutations and exon deletions are most common (5). Procopis et al. (6) reported a mild form of the disorder. This patient had a mutation (c.4085C>T) in the ATP7A gene. Hypotonia, ataxia, and dysmorphic facial features were major clinical findings of this patient.

Lin et al. (7) reported a truncating de novo point mutation in a patient with severe MD who presented with hypotonia, epilepsy, and dysmorphic facial features.

Choi et al. (8) reported the same mutation (p. G1118S [c.3352G>A]) in a 5-month-old male infant who had seizures at the time of diagnosis and was bedridden when he was 3 years old, similar to the present case.

Clinical features, molecular genetic analyses, and disease severity of the patients with MD are shown in Table 1.

The diagnosis was possible at around 9 months in our case and only symptomatic treatment was applied; seizures were controlled after antiepileptic treatment. There have been few reports about the long-term survival in MD. Tchan et al. (9) reported a patient (previously reported by Procopis et al.) who received copper supplementation for >30 years with mild intellectual impairment.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as: Soylu Üstkoyuncu P, Güven AS, Kiraz A, Yılmaz A, Bozdemir ŞE, Gökay S. De Novo Mutation in ATP7A Gene with Severe Menkes Disease. *Erciyes Med J* DOI: 10.5152/etd.2018.0103

Disease variants have been reported in some females and are associated with translocations on X-chromosome. Heterozygous females are thought to be asymptomatic and one-half of the carrier females were shown regions of pili torti. Mutation-targeted molecular testing was performed on the females of the family at risk, and no mutation has been identified in our study. Gu et al. (10) reported that 25% of mothers of the patients with MD were not carriers similar to the present family.

In conclusion, although it is not specific for the disorder, microscopy of the hair allows early diagnosis when the differential diagnosis is broad and/or other tests are not conclusive. Also, serum copper and ceruloplasmin levels are not specific for MD. Therefore, molecular confirmation has to be made for the correct diagnosis. Since it is a fatal disorder, the genetic counseling must be provided to the family in the presence of an index case.

**Informed Consent:** Written informed consent was obtained from the parents of the patient who participated in this study.

**Peer-review:** Externally peer-reviewed.

**Author Contributions:** Conceived and designed the experiments or case: PSÜ, ASG. Performed the experiments or case: AK, AY. Analyzed the data: ŞEB, SG. Wrote the paper: PSÜ.

All authors have read and approved the final manuscript.

**Conflict of Interest:** No conflict of interest was declared by the authors.

**Financial Disclosure:** The authors declared that this study has received no financial support.

## REFERENCES

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as: Soylu Üstkoyuncu P, Güven AS, Kiraz A, Yılmaz A, Bozdemir ŞE, Gökay S. De Novo Mutation in ATP7A Gene with Severe Menkes Disease. *Erciyes Med J* DOI: 10.5152/etd.2018.0103

1. Kaler SG. In: Pagon RA, Adam MP, Ardinger HH, Wallace SE, Amemiya A, Bean LJM, et al. (eds.) GeneReviews® [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2016. 2003 May 9 [updated 2016 Aug 18].
2. Ojha R, Prasad AN. Menkes disease: what a multidisciplinary approach can do. *J Multidiscip Healthc.* 2016; 17(9): 371–85.
3. Rizk T, Mahmoud A, Jamali T, Al-Mubarak S. Menkes Disease Presenting with Epilepsia Partialis Continua. *Case Rep Neurol Med.* 2014; 2014: 525784.
4. Craiu D, Kaler S, Craiu M. Role of optic microscopy for early diagnosis of Menkes disease. *Rom J Morphol Embryol.* 2014; 55(3): 953–56.
5. Tumer Z. An overview and update of ATP7A mutations leading to Menkes disease and occipital horn syndrome. *Hum Mutat.* 2013; 34(3): 417-29.
6. Procopis P, Camakaris J, Danks DM. A mild form of Menkes steely hair syndrome. *J Pediatr.* 1981; 98(1):97-9.
7. Lin YJ, Ho CS, Hsu CH, Lin JL, Chuang CK, Tsai JD, et al. A Truncating De Novo Point Mutation in a Young Infant with Severe Menkes Disease. *Pediatr Neonatol* 2014; 58(1): 1-4.
8. Choi JH, Kim GH, Yoo HW. Identification of novel mutations of the ATP7A gene and prenatal diagnosis of Menkes disease by mutation analysis. *J Genet Med* 2007; 4(1): 38-44.
9. Tchan MC, Wilchen B, Christodoulou J. The mild form of Menkes disease: a 34-year progress report on the original case. *JIMD Rep* 2013; 9: 81–4.
10. Gu YH, Kodama H, Sato E, Mochizuki D, Yanagawa Y, Takayanagi M, et al. Prenatal diagnosis of Menkes disease by genetic analysis and copper measurement. *Brain Dev* 2002; 24(7): 715-8.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as: Soylu Üstkoyuncu P, Güven AS, Kiraz A, Yılmaz A, Bozdemir ŞE, Gökay S. De Novo Mutation in ATP7A Gene with Severe Menkes Disease. *Erciyes Med J* DOI: 10.5152/etd.2018.0103

**Table 1.** Clinical features, molecular genetic analyses, and disease severity of the patients with Menkes Disease

| Author                   | Country      | Age onset | Mutation   | Clinical features  | Disease severity |
|--------------------------|--------------|-----------|------------|--|------------------|
| Procopis et al.,<br>1981 | Australia    | 21 months | c.4085C>T  | Hypotonia, ataxia, dysmorphic facial features  | Mild form        |
| Choi et al.,<br>2003     | Korea        | 5 months  | c.3352G>A  | Seizures   | Severe           |
| Rizk et al.,<br>2014     | Saudi Arabia | 17 months | c.1138G>A  | Epilepsia partialis continua, regression of milestones, repetitive episodes of hypothermia | Severe           |
| Lin et al.,<br>2014      | Taiwan       | 4 months  | c.3502 C>T | Hypotonia, epilepsy, dysmorphic facial features  | Severe           |
| Current report           | Turkey       | 8 months  | c.3352G>A  | Hypotonia, epilepsy, poor feeding  | Severe           |

Figure Legends:


Figure 1. a, b. General view (a) of the patient's hair and pili torti (b)

Figure 2. a-d. Sagittal (a) and coronal (b) and axial (c) T2W images of MRI show diffuse cerebral and cerebellar atrophy and MRA (d) image shows severe intracranial and extracranial vascular tortuosity in both middle cerebral and vertebrobasilar arteries

Figure 3. a-c. Next generation sequencing image (a) shows hemizygous [p. G1118S (c.3352G>A)] mutation in ATP7A gene and the sanger sequencing image (b) shows the confirmation of the patient and the females at risk and the pedigree (c) of the family

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as: Soylu Üstkoyuncu P, Güven AS, Kiraz A, Yılmaz A, Bozdemir ŞE, Gökay S. De Novo Mutation in ATP7A Gene with Severe Menkes Disease. *Erciyes Med J* DOI: 10.5152/etd.2018.0103






This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as: Soylu Üstkoyuncu P, Güven AS, Kiraz A, Yılmaz A, Bozdemir ŞE, Gökay S. De Novo Mutation in ATP7A Gene with Severe Menkes Disease. *Erciyes Med J* DOI: 10.5152/etd.2018.0103

©Copyright 2018 by Erciyes University Faculty of Medicine - Available online at [www.erciyesmedj.com](http://www.erciyesmedj.com)




This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as: Soylu Üstkoyuncu P, Güven AS, Kiraz A, Yılmaz A, Bozdemir ŞE, Gökay S. De Novo Mutation in ATP7A Gene with Severe Menkes Disease. Erciyes Med J DOI: 10.5152/etd.2018.0103

©Copyright 2018 by Erciyes University Faculty of Medicine - Available online at [www.erciyesmedj.com](http://www.erciyesmedj.com)




This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as: Soylu Üstkoyuncu P, Güven AS, Kiraz A, Yılmaz A, Bozdemir ŞE, Gökay S. De Novo Mutation in ATP7A Gene with Severe Menkes Disease. *Erciyes Med J* DOI: 10.5152/etd.2018.0103

©Copyright 2018 by Erciyes University Faculty of Medicine - Available online at [www.erciyesmedj.com](http://www.erciyesmedj.com)





This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as: Soylu Üstkoyuncu P, Güven AS, Kiraz A, Yılmaz A, Bozdemir ŞE, Gökay S. De Novo Mutation in ATP7A Gene with Severe Menkes Disease. Erciyes Med J DOI: 10.5152/etd.2018.0103

©Copyright 2018 by Erciyes University Faculty of Medicine - Available online at [www.erciyesmedj.com](http://www.erciyesmedj.com)



This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as: Soylu Üstkoynucu P, Güven AS, Kiraz A, Yılmaz A, Bozdemir ŞE, Gökay S. De Novo Mutation in ATP7A Gene with Severe Menkes Disease. *Erciyes Med J* DOI: 10.5152/etd.2018.0103

©Copyright 2018 by Erciyes University Faculty of Medicine - Available online at [www.erciyesmedj.com](http://www.erciyesmedj.com)



This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as: Soylu Üstkoyuncu P, Güven AS, Kiraz A, Yılmaz A, Bozdemir ŞE, Gökay S. De Novo Mutation in ATP7A Gene with Severe Menkes Disease. *Erciyes Med J* DOI: 10.5152/etd.2018.0103